


Antimicrobial activity of fluorosubstituted diazines and triazolodiazines.

A.V.Karavaeva, L.A. Kossior, S.V.Vershilov, L.M. Popova

The study of biological activity of fluorine-containing heterocyclic compounds is one of the most important directions of investigations of this class compounds [1-3].

To continue works [4,5] started earlier, we have carried out synthesis [6,7] and evaluation of antimicrobial activity of series of 6-perfluorosubstituted 2-mercapto-4-methylpyrimidines (compounds 1 and 2), 2-perfluorosubstituted 5-methyl-s-triazolo[1,5-a]pyrimidine-7-ones (3-5) and 5-perfluorosubstituted 7-methyl-s-triazolo[1,5-a]pyrimidine (6).



Rf = CF₃ (5); C₆F₁₃ (2,3); CF(CF₃)OC₃F₇ (1,4); CF(CF₃)OCF₂CF(CF₃)OC₃F₇.

Compounds 1-6 were tested in the following cultures: Bac.Cereus ATCC 10702; Ps. Aerogenosa ATCC 900027; E. Coili ATCC 25922 St. and Aureus 209-17.

Table. Antimicrobial activity of fluorosubstituted diazines (1,2) and triazolodiazines (3,6)

Compound	Ds,mm (Cmin,mg/mL)			
	Bac.Cereus	St.Aureus	Ps.Aerogenosa	E.Coli
1.	11 (-)	10(<1.56)	8 (≥1)	0 abs
2.	9 (-)	0 (-)	8 (1)	0 abs
3.	23* (<1.0)	≥30* (<1.0)	8 (-)	0 abs
4.	19* (<1.0)	≥30* (<1.0)	9 (-)	0 abs
5.	- (-)	- (1.56)	- (≥)	0 abs
6.	15* (1.0)	≥30* (<1.0)	9 (-)	0 abs

To determine antimicrobial activity of synthesized compounds 1-6 the following methods were used:

1. Method of diffusion into agar (nutrient medium: MPB, daily culture; microbial load: 10⁹ cells/mL; hole diameter: 6 mm; concentration of the investigated material in the solution: 10 mg/mL).
2. Method of serial dilutions (nutrient medium: MPB, daily culture; microbial load: 10³ cells/mL)

The activity characteristics of the substances, diameter of growth suppression zone of the test-cultures Ds and minimal growth retarding concentrations Cmin are presented in the table.

The check standard is the solvent (ethanol).

Thus, as it follows from the mentioned experimental data, investigated perfluorosubstituted diazines (1,2) and triazolodiazines (3-6) show antimicrobial activity in three cultures except E.Coli. Triazolopyrimidines (3,4 and 6) display distinct antimicrobial activity towards G+ microorganisms, that is obviously connected with the cell shell structure [8].

References

1. Patrik T.b. J.Chem.Soc. 1979.V.56,N4.P.228-230
2. Pat. 5262380 USA.ICI A 021 N 25/32, 43/54.CRJ 5(1), 1995. 50302 P
3. Pat.4904301 USA ICI A 01 N 43/90, C 07 D 487/04 Filed on 19.04.88publ. on 27.02.90. NCI 71.92
4. Siniavskaya, I.M. Kuznetsova, G.G.Nianikova, L.M.Popova, A.I.Trishina, S.V.Vershilov. Articles : Preservation of the environment, issues of ecology and control of product quality. M., NIITEH, 1996. N3, p.7.
5. L.M. Popova, A.Yu.Trishina, S.V.Vershilov, G.G.Nianikova, P.A.Pashkov. Fluorine Notes.1999.N2.
3. L.M.Popova, A.U.Trishina, S.V.Vershilov, A.I Ginak, B.N. Maksimov. J. Fluor. Chem. 1999. V.96. P.51-56.
7. S.V.Vershilov. L.M. Popova. Fluorine Notes.1999.N3

3. Franklin T., Snow G. Biochemistry of antimicrobial action. Moscow; Mir, 1984. 240 p.

Journal content